




AGRISUD INTERNATIONAL



EUROPEAN UNION

កម្មវិធីគាំទ្រការអភិវឌ្ឍន៍សន្តិសុខស្បៀង នៅក្នុងខេត្តបន្ទាយមានជ័យ



ការបណ្តុះផ្សិតចំបើង



ផ្សិតចំបើងអាចបណ្តុះបាននៅក្នុង
រដូវប្រាំង គឺចាប់ពីខែធ្នូ ដល់ខែ
កុម្ភៈ ។ ចំពោះផ្សិតចំបើង គឺជាការ
ពេញនិយមនៅលើទីផ្សារហើយមាន
តំលៃខ្ពស់ទៀតផង ។ ផ្សិតចំបើង
អាចបណ្តុះក្នុងរយៈពេលខ្លីអាចទទួល
ផលបាន ។



វត្តមានដើមដែលអាចយកមកបណ្តុះដ្យូតចំបើងបាន

ជញ្ជាំង ចំបើង ស្លឹកចេកងាប់ ដើមកំប្លោកងាប់



បច្ចេកទេសក្នុងការបណ្តុះដីតំបែង



រៀបចំកន្លែងផ្ទៃរងបណ្តុះដែលមាន
ទទឹង១ ម៉ែត្រ និង បណ្តោយ
ប្រមាណ ៨-១០ ម៉ែត្រ ឬលើសពី
នេះក៏បានដែរ ។

ត្រូវជំរះស្មៅសំអាតនៅលើផ្ទៃរង
អោយអស់ ។



បន្លាបមកបាច Sift lime on
the row to kill germs and
neutralize the soil.



- **Hill up the row. Cultivate the soil.**
- **Dry the cultivated soil from 3 to 5 days.**



- **Roll the rice straw in lump with the same width as row.**
- **Put 1 layer of rolled rice straw on the row. Do it for 3 times.**

- **The more the number of rolled rice straw layers on the row with more mycelium are put, the more the yield is obtained.**
- **Sift well decomposed cowdung on the straw.**
- **Sift mycelium on the row.**





- Cover rice straw on the row and after burn it to increase potassium.



- Water the burned row.



- **Cover the row with plastic rubber to keep heat inside the row. Cover another layer in sheet of thatch or rice straw over the plastic rubber.**

- **Leave the row for 15 days.**
- **Check regularly humidity and heat.**

ការប្រមូលផល (Harvest)



- **Pick the mushroom early morning.**
- **Cover the row immediately after picking to keep heat.**
- **Mushroom can be harvested from 2 to 3 times.**

ស្រមោតក្រណា

